

The detection of extragalactic ^{15}N : Consequences for nitrogen nucleosynthesis and chemical evolution^{1,2}

Yi-nan Chin

Institute of Astronomy and Astrophysics, Academia Sinica, P.O.Box 1-87, Nankang, 11529
Taipei, Taiwan; einmann@asiaa.sinica.edu.tw

Christian Henkel

Max-Planck-Institut für Radioastronomie, Auf dem Hügel 69, D-53121 Bonn, Germany;
p220hen@mpifr-bonn.mpg.de

European Southern Observatory, Casilla 19001, Santiago 19, Chile

Norbert Langer

Institut für Physik, Universität Potsdam, Am Neuen Palais 10, D-14469 Potsdam,
Germany; ntl@astro.physik.uni-potsdam.de

and

Rainer Mauersberger

Steward Observatory, The University of Arizona, Tucson, AZ 85721, U.S.A.;
mauers@as.arizona.edu

Received _____; accepted _____

¹Based on observations with the Swedish-ESO Submillimeter Telescope (SEST) at the European Southern Observatory (ESO), La Silla, Chile

²Please send offprint request to Y.-N. Chin, ASIAA, Taiwan;
einmann@asiaa.sinica.edu.tw

ABSTRACT

Detections of extragalactic ^{15}N are reported from observations of the rare hydrogen cyanide isotope HC^{15}N toward the Large Magellanic Cloud (LMC) and the core of the (post-) starburst galaxy NGC 4945. Accounting for optical depth effects, the LMC data from the massive star-forming region N113 infer a $^{14}\text{N}/^{15}\text{N}$ ratio of 111 ± 17 , about twice the $^{12}\text{C}/^{13}\text{C}$ value. For the LMC star-forming region N159HW and for the central region of NGC 4945, $^{14}\text{N}/^{15}\text{N}$ ratios are also ≈ 100 . The $^{14}\text{N}/^{15}\text{N}$ ratios are smaller than all interstellar nitrogen isotope ratios measured in the disk and center of the Milky Way, strongly supporting the idea that ^{15}N is predominantly synthesized by massive stars. Although this appears to be in contradiction with standard stellar evolution and nucleosynthesis calculations, it supports recent findings of abundant ^{15}N production due to rotationally induced mixing of protons into the helium-burning shells of massive stars.

Subject headings: Nuclear reactions, nucleosynthesis, abundances – ISM: abundances – ISM: molecules – galaxies: abundances – Magellanic Clouds – galaxies: starburst

1. Introduction

Carbon, nitrogen, and oxygen, the ‘CNO-elements’, are among the most abundant species after hydrogen and helium. Being formed by p- and α -capture reactions in the interior of stars, the CNO nuclei are partially released by means of stellar winds, planetary nebula ejecta, and supernova explosions into the interstellar medium (ISM). ^{12}C and ^{16}O and, qualitatively, ^{13}C and ^{17}O nucleosynthesis appears to be understood (e.g., Wilson & Matteucci 1992; Henkel & Mauersberger 1993; Henkel et al. 1994b; Prantzos et al. 1996). The dominant cooking site of the rare nitrogen isotope, ^{15}N , is however not yet known. While ^{14}N is produced in both high and lower mass stars, it is not yet clear whether ^{14}N or ^{15}N is the more ‘primary’ isotope (see Sect. 4).

To obtain new constraints for nitrogen nucleosynthesis, we measured $^{14}\text{N}/^{15}\text{N}$ abundance ratios in so far unexplored molecular environments. The molecular studies by Johansson et al. (1994), Chin et al. (1996, 1997, 1998), and Heikkilä et al. (1997, 1998) demonstrate that it is possible to detect rare molecular species in the Magellanic Clouds. Since such metal poor environments may also characterize cosmologically relevant sources at high redshift, we searched for HC^{15}N in three Magellanic star-forming regions showing prominent $\text{H}^{12}\text{C}^{14}\text{N}$ (hereafter HCN) emission. In view of the low $^{14}\text{N}/^{15}\text{N}$ ratios predicted by Henkel & Mauersberger (1993) and Henkel et al. (1994b) for starbursts, we also searched for HC^{15}N in the southern (post-) starburst galaxy NGC 4945.

2. Observations

The data were taken in September and November 1997 and in January and July 1998 using the 15-m Swedish-ESO Submillimetre Telescope (SEST) at La Silla, Chile. A 3 mm SIS receiver yielded overall system temperatures, including sky noise, of order $T_{\text{sys}} = 250\text{ K}$

on a main beam brightness temperature (T_{MB}) scale. The backend was an acousto-optical spectrometer (AOS). A channel separation of 42 kHz corresponding to 0.14 km s^{-1} at 88 GHz was employed for the observations toward the Magellanic Clouds, while the low resolution spectrometer with a channel separation of 0.7 MHz (or 2.4 km s^{-1}) was used for NGC 4945. The antenna beamwidth was $55''$ at the observed line frequencies taken from Lovas (1992).

The observations were carried out in a dual beam-switching mode (switching frequency 6 Hz) with a beam throw of $11'40''$ in azimuth. All spectral intensities were converted to a T_{MB} scale, correcting for a main beam efficiency of 0.76 (L.B.G. Knee, L.-Å. Nyman, A.R. Tieftrunk, priv. comm.). Calibration was checked by monitoring on Orion KL and NGC 4945 and was found to be consistent between different observation periods within $\pm 10\%$. The pointing accuracy, obtained from measurements of the SiO masers R Dor and W Hya, was better than $10''$.

3. Results

Toward the prominent LMC star-forming regions N113 and N159HW we have detected the $J = 1 - 0$ emission lines of hydrogen cyanide (HCN) and its rare isotopic species H^{13}CN and HC^{15}N . Although they were also detected toward the core of the (post-) starburst galaxy NGC 4945 (e. g., Koornneef 1993), only HCN was seen in the SMC star-forming region LIRS 36 (see Chin et al. 1998). Spectra and line parameters obtained from Gaussian fits are displayed in Fig. 1 and Table 1. Observed line intensity ratios, derived opacities, and isotopic abundance ratios ($^{12}\text{C}/^{13}\text{C}$ and $^{14}\text{N}/^{15}\text{N}$) are given in Table 2.

Apart from optical depth effects, integrated line intensity ratios should be identical to isotope ratios within the observational errors, since rotational constants of the various HCN isotopomers are similar. A fractionation of nitrogen isotopes via charge exchange reactions,

as expected in the case of carbon (see Watson et al. 1976), must be negligible, because the nitrogen ionization potential is higher than that of hydrogen; N^+ abundances are small in molecular clouds. For the two LMC star-forming regions and for NGC 4945 we then find $^{14}\text{N}/^{15}\text{N} \approx 100$. In the case of N113, the $^{14}\text{N}/^{15}\text{N}$ ratio does not depend on an assumption on $^{12}\text{C}/^{13}\text{C}$. Instead, $^{14}\text{N}/^{15}\text{N}$ is *directly* derived from the $\text{HCN}/\text{HC}^{15}\text{N}$ integrated line temperature ratio and from a fit to the HCN hyperfine components (see Table 2). Toward N159HW, the HCN main line also appears to be almost optically thin but a larger intrinsic linewidth makes the fit less convincing.

4. Discussion

Observations indicate that ^{14}N production involves both, a primary and a secondary component (e.g., Matteucci 1986; Vila-Costas & Edmunds 1993; van Zee et al. 1998), which is currently understood in terms of primary hydrostatic ^{14}N production in massive low metallicity stars (cf. Laird 1985, Timmes et al. 1995), while the secondary production of ^{14}N through the CNO cycle is dominating for $Z \gtrsim Z_{\odot}/100$ (cf. Timmes et al. 1995; van den Hoek & Groenewegen 1997).

^{15}N is destroyed during hydrostatic hydrogen-burning (e.g., Wannier et al. 1991; El Eid 1994; Langer & Henkel 1995) and is thus thought to be a product of explosive hydrogen nucleosynthesis. It is synthesized in novae as a mostly primary nucleus (e.g., José & Hernanz 1998). However, current models produce insufficient amounts to account for the galactic ^{15}N abundance (Kovetz & Prialnik 1997). Alternatively, it may be produced in Type II supernovae through neutrino spallation of primary ^{16}O (Woosley & Weaver 1995, Timmes et al. 1995; see also Audouze et al. 1977). A small contribution from Type Ia supernovae is also possible (Clayton et al. 1997; although see Nomoto et al. 1984). In the first case (novae), there is a time delay between star formation and the occurrence of

^{15}N enriched ejecta, leading to initially high $^{14}\text{N}/^{15}\text{N}$ ratios that gradually decrease with time (for an illustration, see e. g. Fig. 4 of Güsten & Ungerechts 1985). In the second case (massive stars), rapid injection of ^{15}N into the ISM leads to initially small $^{14}\text{N}/^{15}\text{N}$ ratios that gradually increase with time when the ^{14}N contribution from low-mass stars becomes significant.

Analyzing $^{14}\text{N}/^{15}\text{N}$ data from HCN and NH_3 , it soon became obvious that local ISM and galactic center $^{14}\text{N}/^{15}\text{N}$ ratios are moderately and strongly enhanced relative to the solar system value. Assuming that the solar system, the local ISM, and the galactic center region form a sequence of increasing degree of nuclear processing (the solar system was assumed to reflect the composition of the outer Galaxy at a time almost five billion years ago) led to the conclusion that ^{15}N is predominantly arising from massive stellar progenitors (e. g., Audouze et al. 1977; Güsten & Ungerechts 1985). In this case the more secondary ^{14}N nucleus becomes more abundant relative to ^{15}N with higher degree of nuclear processing, yielding $^{14}\text{N}/^{15}\text{N}$ ratios of 270, 300 – 400, and 500 – 1000 in the solar system, the local ISM, and the galactic center region, respectively.

Two complications led to the present ‘nitrogen puzzle’: Firstly, the Sun is more metal rich than the local ISM (e. g., Russell & Dopita 1992; Cameron 1993; Cunha & Lambert 1994; Meyer 1997), in spite of its age. Secondly, Dahmen et al. (1995) found a galactic disk gradient with $^{14}\text{N}/^{15}\text{N}$ ratios (from HCN) increasing with galactocentric distance. The first complication is easily resolved within the framework of primary ^{15}N production, if we note that the Sun is particularly enriched with nuclei processed in massive stars. This leads to enhanced $^{18}\text{O}/^{17}\text{O}$ ratios (Henkel & Mauersberger 1993; Henkel et al. 1994b) and to reduced $^{14}\text{N}/^{15}\text{N}$ ratios relative to the local ISM. The $^{14}\text{N}/^{15}\text{N}$ gradient is a more severe problem. The gradient implies that either the initial mass function (IMF) is biased in favor of high mass stars in the inner galactic disk (there is no observational support for such a

bias) or that infall of halo gas or of cannibalized dwarf galaxies have altered interstellar abundances. Most galactic $^{14}\text{N}/^{15}\text{N}$ values are determined from $^{14}\text{N}/^{15}\text{N} : ^{12}\text{C}/^{13}\text{C}$ double isotope ratios. Carbon fractionation may lead to small deviations between $\text{HCN}/\text{H}^{13}\text{CN}$ and $^{12}\text{C}/^{13}\text{C}$ abundance ratios (see Langer et al. 1984, 1989; Wilson & Matteucci 1992; Wilson & Rood 1994), but this should not alter the sign of a gradient. Assuming instead delayed injection of ^{15}N via novae into the ISM (e. g., Dahmen et al. 1995) fails to account for the large measured $^{14}\text{N}/^{15}\text{N}$ ratios in the galactic center region with its high number of potential nova candidates (see Shafter 1997).

Solving the nitrogen puzzle requires the determination of small (≈ 100) $^{14}\text{N}/^{15}\text{N}$ ratios in a (post-) starburst environment, that is strongly influenced by the ejecta from massive stars (Henkel & Mauersberger 1993; Henkel et al. 1994b). Only a lower $^{14}\text{N}/^{15}\text{N}$ limit of order 100 was obtained toward M 82 (Henkel et al. 1998). Our $^{14}\text{N}/^{15}\text{N}$ ratio from NGC 4945 strongly suggests that the bulk of ^{15}N is ejected by massive stars.

The ISM of the LMC is less processed than the solar system and the local ISM and should also be characterized by overabundances of nuclei ejected from massive stars. $[\text{C}/\text{O}]_{\text{LMC}} \sim [\text{N}/\text{O}]_{\text{LMC}} \sim -0.3$ is consistent with this hypothesis (e. g., Westerlund 1990). Thus, if ^{15}N is released by massive stars, an ‘overabundance’ of ^{15}N relative to ^{14}N is expected; in the LMC, $^{14}\text{N}/^{15}\text{N}$ should be significantly smaller than ratios measured in the solar system and the local ISM.

Our observational results from the LMC and NGC 4945 are thus providing strong support for ^{15}N being predominantly ejected by Type II supernovae. This finding is not consistent with the sign of the galactic disk $^{14}\text{N}/^{15}\text{N}$ gradient (Dahmen et al. 1995). It also seems to be in conflict with numerical calculations of massive star nucleosynthesis (e. g., Weaver & Woosley 1993; Woosley & Weaver 1995). While without the contribution of neutrino-induced nucleosynthesis to the production of ^{15}N , massive stars would destroy

rather than produce this isotope (cf. Weaver & Woosley 1993), Woosley & Weaver (1995) find the neutrino production with the (uncertain) currently predicted neutrino scattering cross sections not sufficient to explain the solar system abundance of ^{15}N (cf. also Timmes et al. 1995). However, recent massive star models which take the effects of rotation on the stellar structure and nucleosynthesis into account (Heger et al. 1997, 1999, Langer et al. 1999) indicate the possibility of abundant hydrostatic production of ^{15}N in case of mixing between the hydrogen-burning and the helium-burning shell, a mechanism already discussed by Jorissen & Arnould (1989). Consequently, although there are considerable uncertainties and weakly restricted parameters in the rotating massive star models which need to be explored in the near future, massive stars should be considered as an important source of ^{15}N in galaxies, particularly so after the observational facts reported in the present paper.

We thank J.H. Black and U. Ott for useful discussions and criticism. YNC thanks for financial support through National Science Council of Taiwan grant 86-2112-M001-032. RM acknowledges support from NATO grant SA.5-2-05 (CRG. 960086) 318/96, and NL from the Deutsche Forschungsgemeinschaft through grants La 587/15-1 and 16-1.

REFERENCES

- Audouze, J., Lequeux, J., Rocca-Volmerange, & B., Vigroux, L. 1977, CNO Isotopes in Astrophysics, ed. J. Audouze, D. Reidel Publishing Company, Dordrecht, p155
- Cameron, A.G.W. 1993, Protostars & Planets III, eds. E.H. Levy, J.I. Lunine, The University of Arizona Press, Tucson, p47
- Chin, Y.-N., Henkel, C., Millar, T.J., Whiteoak, J.B., & Mauersberger, R., 1996, A&A, 312, L33
- Chin, Y.-N., Henkel, C., Whiteoak, J.B., et al. 1997, A&A, 317, 548
- Chin, Y.-N., Henkel, C., Millar, T.J., Whiteoak, J.B., & Marx-Zimmer, M. 1998, A&A, 330, 901
- Clayton, D.D., Arnett, D., Kane, J., & Meyer, B.S. 1997, ApJ, 486, 824
- Cunha, K., & Lambert, D.L. 1994, ApJ, 426, 170
- Dahmen, G., Wilson, T.L., & Matteucci, F. 1995, A&A, 295, 194
- El Eid, M.F. 1994, A&A, 285, 915
- Güsten, R., & Ungerechts, H. 1985, A&A, 145, 241
- Heger, A., Jeannin, L., Langer, N., & Baraffe, I. 1997, A&A, 327, 224
- Heger, A., Woosley, S.E., & Langer, N. 1999, ApJ, in preparation
- Heikkilä, A., Johansson, L.E.B., & Olofsson, H. 1997, A&A, 319, L21
- Heikkilä, A., Johansson, L.E.B., & Olofsson, H. 1998, A&A, 332, 493
- Henkel, C., & Mauersberger, R. 1993, A&A, 274, 730

- Henkel, C., Whiteoak, J.B., & Mauersberger, R. 1994a, *A&A*, 284, 17
- Henkel, C., Wilson, T.L., Langer, N., Chin, Y.-N., & Mauersberger, R. 1994b, *The Structure and Content of Molecular Clouds*, eds. T.L. Wilson, K.J. Johnston, Springer Verlag, Berlin, p72
- Henkel, C., Chin, Y.-N., Mauersberger, R., & Whiteoak, J.B. 1998, *A&A*, 329, 443
- Johansson, L.E.B., Olofsson, H., Hjalmarson, Å, Gredel, R., & Black, J.H., 1994, *A&A*, 291, 89
- Jorissen, A., & Arnould, M. 1989, *A&A*, 221, 161
- José, J., & Hernanz, M. 1998, *ApJ*, 494, 680
- Koornneef, J. 1993, *ApJ*, 403, 581
- Kovetz, A., & Prialnik, D. 1997, *ApJ*, 477, 356
- Laird, J.B. 1985, *ApJ*, 289, 556
- Langer, N., & Henkel, C. 1995, *Sp. Sci. Rev.*, 74, 343
- Langer, N., Heger, A., & García-Segura, G. 1999, *Rev. Modern Astron.*, Vol. 11, in press
- Langer, W.D., & Graedel, T.E. 1989, *ApJS*, 69, 241
- Langer, W.D., Graedel, T.E., Frerking, M.A., & Armentrout, P.B. 1984, *ApJ*, 277, 581
- Lovas, F.J. 1992, *J. Phys. Chem. Ref. Data*, 21, 181
- Matteucci, F. 1986, *MNRAS*, 221, 911
- Meyer, D.M. 1997, *IAU Symp. 178, Molecules in Astrophysics: Probes and Processes*, ed. E.F. van Dishoeck, Kluwer Academic Publishers, Dordrecht, p407

- Nomoto, K., Thielemann, F.-K., & Yokoi, K. 1984, *ApJ*, 286, 644
- Prantzos, N., Aubert, O., & Audouze, J. 1996, *A&A*, 309, 760
- Russell, S.C., & Dopita, M.A. 1992, *ApJ*, 384, 508
- Shafter, A.W. 1997, *ApJ*, 487, 226
- Timmes, F.X., Woosley, S.E., & Weaver, T.A. 1995, *ApJS*, 98, 617
- van den Hoek, L.B., & Groenewegen, M.A.T. 1997, *A&AS*, 123, 305
- van Zee, L., Salzer, J.J., & Haynes, M.P. 1998, *ApJ*, 497, L1
- Vila-Costas, M.B., & Edmunds, M.G. 1993, *MNRAS*, 265, 199
- Wannier, P.G., Linke, R.A., & Penzias, A.A. 1981, *ApJ*, 247, 522
- Wannier, P.G., Andersson, B.-G., Olofsson, H., Ukita, N., & Young, K. 1991, *ApJ*, 380, 593
- Watson, W.D., Anicich, V.G., & Huntress, W.T., Jr. 1976, *ApJ*, 205, L165
- Weaver, T.A., & Woosley, S.E. 1993, *Phys. Rep.*, 227, 65
- Westerlund, B.E. 1990, *A&AR*, 2, 29
- Wilson, T.L., & Matteucci, F. 1992, *A&AR*, 4, 1
- Wilson, T.L., & Rood, R.T. 1994, *ARAA*, 32, 191
- Woosley, S.E., & Weaver, T.A. 1995, *ApJS*, 101, 181

Fig. 1.— Spectra of HCN, H^{13}CN , and HC^{15}N ; for coordinates, see Chin et al. (1997) and Henkel et al. (1994a)

Table 1. Molecular lines parameters

Molecule & Transition		Frequency [GHz]	T_{MB} [mK]	r.m.s. ^a [mK]	v_{LSR} [km s ⁻¹]	$\Delta v_{1/2}$ [km s ⁻¹]	$\int T_{\text{MB}} dv$ [K km s ⁻¹]
N113							
HCN ^b	$J=1-0$ {	$F=1-1$	88.630416	158			0.809 ± 0.057
		$F=2-1$	88.631847	246	11	235.0	1.26 ± 0.06
		$F=0-1$	88.633936	68			0.346 ± 0.057
H ¹³ CN ^c	$J=1-0$	86.340184	7.3	3	235.4	8.3	0.065 ± 0.004
HC ¹⁵ N ^c	$J=1-0$	86.054961	7.2	3	235.2	4.6	0.035 ± 0.004
N159HW							
HCN ^b	$J=1-0$ {	$F=1-1$	88.630416	65			0.425 ± 0.087
		$F=2-1$	88.631847	156	13	237.7	1.03 ± 0.09
		$F=0-1$	88.633936	43			0.283 ± 0.087
H ¹³ CN ^c	$J=1-0$	86.340184	5.9	5	239.4	12.0	0.076 ± 0.009
HC ¹⁵ N ^c	$J=1-0$	86.054961	3.2	3	237.2	10.4	0.035 ± 0.005
LIRS 36							
HCN ^b	$J=1-0$ {	$F=1-1$	88.630416	12			0.032 ± 0.018
		$F=2-1$	88.631847	25	7	126.4	0.066 ± 0.019
		$F=0-1$	88.633936	4			0.011 ± 0.018
H ¹³ CN ^c	$J=1-0$	86.340184	< 5	6	—	—	< 0.020 ^d
HC ¹⁵ N ^c	$J=1-0$	86.054961	< 5	6	—	—	< 0.024 ^d
NGC 4945							
HCN	$J=1-0$	88.631847	67	11	550	300	21.3 ± 0.1
H ¹³ CN	$J=1-0$	86.340184	3.0	1	550	300	0.959 ± 0.013
HC ¹⁵ N	$J=1-0$	86.054961	1.8	1	540	250	0.464 ± 0.012

^a1 σ noise level of a single channel width of 0.14 km s⁻¹ (for the Magellanic Clouds) or 2.4 km s⁻¹ (for NGC 4945) on a T_{MB} scale.

^bThe three HCN hyperfine transitions ($F=1-1$, $F=2-1$, $F=0-1$) have been resolved by a Gaussian fit. While T_{MB} values and integrated line intensities are given for each component, the velocity refers to the main component.

^cThe hyperfine components of H¹³CN and HC¹⁵N cannot be resolved.

^dFor undetected lines, 3 σ values are given.

Table 2. Integrated line intensity ratios ‘ R ’ of HCN isotopic species, integrated line intensity ratios ‘ R_{02} ’ and ‘ R_{12} ’ of HCN hyperfine transitions with opacity τ_0 , and isotopic abundance ratios.

Source	$R\left(\frac{\text{HCN}}{\text{H}^{13}\text{CN}}\right)$	$R\left(\frac{\text{HCN}}{\text{HC}^{15}\text{N}}\right)$	$R\left(\frac{\text{H}^{13}\text{CN}}{\text{HC}^{15}\text{N}}\right)$	R_{12}^{a}	R_{02}^{a}	τ_0^{a}	$^{12}\text{C}/^{13}\text{C}$	$^{14}\text{N}/^{15}\text{N}$
N113	37 ± 5	69 ± 13	1.9 ± 0.3	0.64 ± 0.07	0.27 ± 0.06	0.122	60 ± 6	111 ± 17
N159HW	23 ± 6	49 ± 15	2.1 ± 0.6	0.41 ± 0.12	0.27 ± 0.11	< 1	$\approx 50^{\text{b}}$	$\approx 100^{\text{c}}$
LIRS 36	> 4.5	> 5.5	—	0.49 ± 0.41	0.17 ± 0.32	—	—	—
NGC 4945	22 ± 1	46 ± 1	2.1 ± 0.1	—	—	< 1	$\approx 50^{\text{b}}$	$\approx 100^{\text{c}}$

^aThe opacity of the HCN $J=1-0$ transition can be derived from the line intensity ratios of the $J=1-0$ hyperfine components. The degeneracies of the HCN $J = 1$, $F = 0, 1, 2$ states are 1, 3, and 5. If τ_0 , τ_1 , τ_2 denote the optical depths of the transitions from these three states to the ground state, respectively, we find under conditions of Local Thermodynamical Equilibrium (LTE) $\tau_1 = 3\tau_0$ and $\tau_2 = 5\tau_0$. The intensity ratios of the hyperfine transitions are then $R_{12} = \frac{I(F=1-1)}{I(F=2-1)} = \frac{1 - e^{-3\tau_0}}{1 - e^{-5\tau_0}}$ and $R_{02} = \frac{I(F=0-1)}{I(F=2-1)} = \frac{1 - e^{-\tau_0}}{1 - e^{-5\tau_0}}$. For NGC 4945, τ_0 was estimated from $^{12}\text{C}/^{13}\text{C}$ (Henkel et al. 1994a) and the observed line intensity ratio (see Table 1).

^bFor the $^{12}\text{C}/^{13}\text{C}$ ratio in N159, see Johansson et al. (1984); for NGC 4945, see Henkel et al. (1994a).

^cFor NGC 4945, lines are too broad to see individual hyperfine components in the HCN line, but the $^{14}\text{N}/^{15}\text{N}$ ratio can be estimated by $^{14}\text{N}/^{15}\text{N} = \frac{\text{H}^{13}\text{CN}}{\text{HC}^{15}\text{N}} \times ^{12}\text{C}/^{13}\text{C}$. This can also be used to confirm the $^{14}\text{N}/^{15}\text{N}$ ratio determined in N159HW.

